

# BR30310 Advanced Equine Nutrition

View Online



---

[1]

Abdouli, H. and Attia, S.B. 2007. Evaluation of a two-stage in vitro technique for estimating digestibility of equine feeds using horse faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 132, 1-2 (Jan. 2007), 155-162.  
DOI:<https://doi.org/10.1016/j.anifeedsci.2006.03.005>.

[2]

Dougal, K. et al. 2012. A comparison of the microbiome and the metabolome of different regions of the equine hindgut. *FEMS Microbiology Ecology*. 82, 3 (Dec. 2012), 642-652.  
DOI:<https://doi.org/10.1111/j.1574-6941.2012.01441.x>.

[3]

Frape, D. 2010. *Equine nutrition and feeding*. Wiley-Blackwell.

[4]

Lowman, R.S. et al. 1999. Evaluation of an in vitro batch culture technique for estimating the in vivo digestibility and digestible energy content of equine feeds using equine faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 80, 1 (Jul. 1999), 11-27. DOI:[https://doi.org/10.1016/S0377-8401\(99\)00039-5](https://doi.org/10.1016/S0377-8401(99)00039-5).

[5]

McDonald, P. 2011. *Animal nutrition*. Pearson.

[6]

National Research Council (U.S.) 2007. Nutrient requirements of horses. National Academies Press.