

BR30310 Advanced Equine Nutrition

View Online



Abdouli, H., and S. Ben Attia. 'Evaluation of a Two-Stage in Vitro Technique for Estimating Digestibility of Equine Feeds Using Horse Faeces as the Source of Microbial Inoculum'. *Animal Feed Science and Technology* 132.1-2 (2007): 155-162. Web.

Dougal, Kirsty et al. 'A Comparison of the Microbiome and the Metabolome of Different Regions of the Equine Hindgut'. *FEMS Microbiology Ecology* 82.3 (2012): 642-652. Web.

Frape, David. *Equine Nutrition and Feeding*. 4th ed. Chichester: Wiley-Blackwell, 2010. Print.

Lowman, R.S et al. 'Evaluation of an in Vitro Batch Culture Technique for Estimating the in Vivo Digestibility and Digestible Energy Content of Equine Feeds Using Equine Faeces as the Source of Microbial Inoculum'. *Animal Feed Science and Technology* 80.1 (1999): 11-27. Web.

McDonald, Peter. *Animal Nutrition*. 7th ed. Harlow, England: Pearson, 2011. Web.
<http://eu.alma.exlibrisgroup.com/view/action/uresolver.do?operation=resolveService&package_service_id=3037249640002418&institutionId=2418&customerId=2415>.

National Research Council (U.S.). *Nutrient Requirements of Horses*. 6th rev. ed. Washington, D.C.: National Academies Press, 2007. Print.