

# BR30310 Advanced Equine Nutrition

View Online



- 
1.  
Frape, D.: Equine nutrition and feeding. Wiley-Blackwell, Chichester (2010).
  
  2.  
Abdouli, H., Attia, S.B.: Evaluation of a two-stage in vitro technique for estimating digestibility of equine feeds using horse faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 132, 155–162 (2007).  
<https://doi.org/10.1016/j.anifeedsci.2006.03.005>.
  
  3.  
Lowman, R.S., Theodorou, M.K., Hyslop, J.J., Dhanoa, M.S., Cuddeford, D.: Evulation of an in vitro batch culture technique for estimating the in vivo digestibility and digestible energy content of equine feeds using equine faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 80, 11–27 (1999).  
[https://doi.org/10.1016/S0377-8401\(99\)00039-5](https://doi.org/10.1016/S0377-8401(99)00039-5).
  
  4.  
McDonald, P.: Animal nutrition. Pearson, Harlow, England (2011).
  
  5.  
National Research Council (U.S.): Nutrient requirements of horses. National Academies Press, Washington, D.C. (2007).
  
  - 6.

Dougal, K., Harris, P.A., Edwards, A., Pachebat, J.A., Blackmore, T.M., Worgan, H.J., Newbold, C.J.: A comparison of the microbiome and the metabolome of different regions of the equine hindgut. *FEMS Microbiology Ecology*. 82, 642–652 (2012).  
<https://doi.org/10.1111/j.1574-6941.2012.01441.x>.