

BR30310 Advanced Equine Nutrition

View Online



Abdouli, H., & Attia, S. B. (2007). Evaluation of a two-stage in vitro technique for estimating digestibility of equine feeds using horse faeces as the source of microbial inoculum. *Animal Feed Science and Technology*, 132(1-2), 155-162.

<https://doi.org/10.1016/j.anifeedsci.2006.03.005>

Dougal, K., Harris, P. A., Edwards, A., Pachebat, J. A., Blackmore, T. M., Worgan, H. J., & Newbold, C. J. (2012). A comparison of the microbiome and the metabolome of different regions of the equine hindgut. *FEMS Microbiology Ecology*, 82(3), 642-652.

<https://doi.org/10.1111/j.1574-6941.2012.01441.x>

Frape, D. (2010). *Equine nutrition and feeding* (4th ed). Wiley-Blackwell.

Lowman, R. S., Theodorou, M. K., Hyslop, J. J., Dhanoa, M. S., & Cuddeford, D. (1999). Evaluation of an in vitro batch culture technique for estimating the in vivo digestibility and digestible energy content of equine feeds using equine faeces as the source of microbial inoculum. *Animal Feed Science and Technology*, 80(1), 11-27.

[https://doi.org/10.1016/S0377-8401\(99\)00039-5](https://doi.org/10.1016/S0377-8401(99)00039-5)

McDonald, P. (2011). *Animal nutrition* (7th ed). Pearson.

http://eu.alma.exlibrisgroup.com/view/action/uresolver.do?operation=resolveService&package_service_id=3037249640002418&institutionId=2418&customerId=2415

National Research Council (U.S.). (2007). *Nutrient requirements of horses* (6th rev. ed). National Academies Press.