

BR30310 Advanced Equine Nutrition

[View Online](#)

[1]

D. Frape, *Equine nutrition and feeding*, 4th ed. Chichester: Wiley-Blackwell, 2010.

[2]

H. Abdouli and S. B. Attia, 'Evaluation of a two-stage in vitro technique for estimating digestibility of equine feeds using horse faeces as the source of microbial inoculum', *Animal Feed Science and Technology*, vol. 132, no. 1-2, pp. 155-162, Jan. 2007, doi: 10.1016/j.anifeedsci.2006.03.005.

[3]

R. S. Lowman, M. K. Theodorou, J. J. Hyslop, M. S. Dhanoa, and D. Cuddeford, 'Evaluation of an in vitro batch culture technique for estimating the in vivo digestibility and digestible energy content of equine feeds using equine faeces as the source of microbial inoculum', *Animal Feed Science and Technology*, vol. 80, no. 1, pp. 11-27, Jul. 1999, doi: 10.1016/S0377-8401(99)00039-5.

[4]

P. McDonald, *Animal nutrition*, 7th ed. Harlow, England: Pearson, 2011. [Online]. Available: http://eu.alma.exlibrisgroup.com/view/action/uresolver.do?operation=resolveService&package_service_id=3037249640002418&institutionId=2418&customerId=2415

[5]

National Research Council (U.S.), *Nutrient requirements of horses*, 6th rev. ed. Washington, D.C.: National Academies Press, 2007.

[6]

K. Dougal et al., 'A comparison of the microbiome and the metabolome of different regions of the equine hindgut', FEMS Microbiology Ecology, vol. 82, no. 3, pp. 642–652, Dec. 2012, doi: 10.1111/j.1574-6941.2012.01441.x.