

BR30310 Advanced Equine Nutrition

View Online



-
1.
Frape, D.: Equine nutrition and feeding. Wiley-Blackwell, Chichester (2010).

 2.
Abdouli, H., Attia, S.B.: Evaluation of a two-stage in vitro technique for estimating digestibility of equine feeds using horse faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 132, 155–162 (2007).
<https://doi.org/10.1016/j.anifeedsci.2006.03.005>.

 3.
Lowman, R.S., Theodorou, M.K., Hyslop, J.J., Dhanoa, M.S., Cuddeford, D.: Evaluation of an in vitro batch culture technique for estimating the in vivo digestibility and digestible energy content of equine feeds using equine faeces as the source of microbial inoculum. *Animal Feed Science and Technology*. 80, 11–27 (1999).
[https://doi.org/10.1016/S0377-8401\(99\)00039-5](https://doi.org/10.1016/S0377-8401(99)00039-5).

 4.
McDonald, P.: *Animal nutrition*. Pearson, Harlow, England (2011).

 5.
National Research Council (U.S.): *Nutrient requirements of horses*. National Academies Press, Washington, D.C. (2007).

 - 6.

Dougal, K., Harris, P.A., Edwards, A., Pachebat, J.A., Blackmore, T.M., Worgan, H.J., Newbold, C.J.: A comparison of the microbiome and the metabolome of different regions of the equine hindgut. *FEMS Microbiology Ecology*. 82, 642–652 (2012).
<https://doi.org/10.1111/j.1574-6941.2012.01441.x>.